

## **REGENERATIVE MEDICINE**

Regenerative Medicine techniques harness the body's innate healing mechanisms to accelerate healing and stimulate tissue regeneration. Our techniques use donor or patient-derived bioactive products designed to promote innate healing mechanisms.

At Hudson Medical, we understand the immense potential of Regenerative Medicine and are dedicated to providing you with the most advanced treatment options that target your specific needs. We use minimally-invasive techniques to treat various tissue injuries by stimulating the regeneration of damaged tissue. These regenerative therapies are designed to repair damaged tissues to reduce the need for more invasive procedures. As a research facility for ongoing clinical trials, we are able to offer cutting-edge regenerative medicine treatment options that are performed by our team of highly experienced physicians. We perform a range of regenerative therapeutic interventions and infusions aimed to treat myriad conditions such as musculoskeletal injuries, degenerative spinal conditions, sports-related injuries, long-term COVID-19-related symptoms, and age-related conditions.

The regenerative therapies we offer at Hudson Medical are appropriate for individuals who struggle to find symptom relief from traditional treatments. Depending on your needs, we implement an individualized treatment plan focused on stimulating tissue regeneration that is tailored to your specific condition.

We now offer Platelet-Rich Plasma Therapy, Platelet-Rich Fibrin Therapy, Bone Marrow Aspirate Concentrate Therapy, Wharton's Jelly, and Extracellular Growth Factors (Exosomes). Hudson Medical is the first practice in the US to publish safety data on the usage of Exosomes in spinal based applications, and will continue to lead research in this field

### **PLATELET RICH PLASMA THERAPY**

Platelet-rich plasma (PRP) Therapy is a minimally-invasive injection that aims to promote the initiation and progression of the body's natural healing mechanisms to accelerate tissue regeneration. Since the 1980s, PRP Therapy has been used in many areas of medicine and dentistry to promote wound healing. In recent years, PRP Therapy has been shown to be safe and effective in the treatment of musculoskeletal injuries, androgenetic alopecia and erectile dysfunction. PRP is derived from a patient's own blood sample and contains high levels of platelets and growth factors known to stimulate tissue healing and reduce inflammation. By directly injecting PRP into sites of tissue injury, this procedure can provide targeted and effective symptom relief by reducing pain and promoting healing. At Hudson Medical, we offer PRP Therapy as an alternative treatment option for individuals with osteoarthritis, tendinopathy, androgenetic alopecia, erectile dysfunction, and chronic back pain due to spinal radiculopathy, spinal spondylosis or sacroiliac joint dysfunction. PRP Therapy is a safe and effective procedure that harnesses innate healing processes to facilitate tissue repair. To learn more about PRP Therapy, please click here [→](#)

### **PLATELET RICH FIBRIN THERAPY**

Platelet-rich fibrin (PRF) Therapy is a minimally-invasive injection used to enhance the regenerative properties of tissue that becomes damaged due to musculoskeletal conditions such as muscle strain injuries, cartilage injuries, and osteoarthritis. PRF contains high levels of cytokines, adhesive proteins, and growth factors that promote tissue regeneration and reduce healing times in such musculoskeletal conditions. Similarly to PRP, PRF product is obtained through a simple blood draw and is injected directly into the site of tissue injury. The unique blend of bioactive substances found in PRF allow for an extended release of these factors after the PRF is injected. At Hudson Medical, we offer PRF Therapy to stimulate regeneration of surrounding healthy tissue in patients with persistent symptoms caused by musculoskeletal conditions. PRF Therapy can provide significant pain relief while simultaneously activating innate healing processes associated with tissue regeneration. To learn more about PRF Therapy, please click here [→](#)

### **WHARTON'S JELLY**

Recent developments in stem cell research have demonstrated the regenerative power of Wharton's Jelly in the treatment of intervertebral disc degeneration, osteoarthritis and chronic tendinopathy. Wharton's Jelly is derived from human umbilical cord tissue and contains high levels of mesenchymal stem cells (MSCs), which have the ability to differentiate into specialized cell types and therefore have profound regenerative potential. Wharton's Jelly also contains a number of other substances such as growth factors, cytokines, hyaluronic acid, and exosomes that can enhance wound healing and promote tissue regeneration. At Hudson Medical, we offer Wharton's Jelly as an alternative treatment option for patients with various orthopedic conditions such as osteoarthritis and intervertebral disc degeneration. Our physicians inject Wharton's Jelly into injury sites to stimulate tissue remodeling, growth and regeneration. To learn more about Wharton's Jelly, please click here [→](#)

### **BONE MARROW ASPIRATE CONCENTRATE**

Bone Marrow Aspirate Concentrate (BMAC) Therapy uses mesenchymal stem cells (MSCs) from patients' bone marrow to stimulate tissue regeneration for a wide range of clinical applications. MSCs are a self-renewing cell type that can transform into different specialized cell types by interacting with signaling molecules implicated in different physiological processes. BMAC Therapy is a minimally-invasive procedure that involves the harvesting of a small amount of a patient's bone marrow and subsequent injection of BMAC product directly into injury sites. At Hudson Medical, we use BMAC Therapy to treat musculoskeletal conditions such as osteoarthritis, tendinopathy and chronic back pain because the MSCs, growth factors, and cytokines found in BMAC products activate tissue repair mechanisms and provide patients with significant pain and symptom relief. BMAC Therapy is a safe and effective treatment that targets physiological causes of musculoskeletal conditions to promote healing and tissue regeneration. To learn more about BMAC Therapy, please click here [→](#)

### **EXOSOME THERAPY**

Exosomes are small vesicles secreted by many cell types that carry diverse cargo to target cells to elicit a cargo-specific response. Exosomes derived from umbilical cord, amniotic fluid or mesenchymal stem cells carry cargo that can alter gene expression of target cells, promoting processes that favor regeneration and healing. As exosomes are able to modify cellular communication and can therefore alter tissue microenvironments, Exosome Therapy has extensive clinical applications for regenerative medicine. The exosome products used in Exosome Therapy are derived from donor stem cells and can be either directly injected into injury sites or administered intravenously depending on the patient's condition. Conditions such as osteoarthritis and intervertebral disc degeneration may benefit from direct Exosome Therapy injection to promote tissue regeneration at damaged tissue, whereas Exosome Therapy intravenous infusions may be better indicated in the treatment plans of aging-related conditions, COVID-19-induced severe acute respiratory distress syndrome, long-term COVID-19 symptoms, and chronic pain conditions. Exosome Therapy stimulates tissue regeneration and cellular rejuvenation by harnessing the intrinsic ability of exosomal cargo to alter cellular communication and gene expression. To learn more about Exosome Therapy, please click here [→](#)

### **WHARTON'S JELLY**

In recent years, regenerative medicine therapies using biologic applications, such as stem cell products and platelet-rich plasma treatments, have gained the attention of many researchers and clinicians due to encouraging experimental evidence. Of the various stem cells formed by the body, mesenchymal stem cells (MSCs), also known as stromal cells, are arising as a promising therapeutic application for musculoskeletal injuries and conditions such as osteoarthritis and intervertebral disc degeneration. Stem cells are cells which have the ability to transform into specialized cell types through interactions with signaling factors called growth factors and cytokines (1). The exact signaling molecules and their associated mechanisms of action are not fully understood, but current research suggests that these signaling factors induce stem cells to divide, differentiate, and proliferate into specialized cell types (1). The regenerative properties of stem cells provides an exciting opportunity to expand the available treatment options for complications associated with organ transplantation, diabetes, neurological conditions, cardiovascular diseases, and bone and cartilage conditions (2). Recent research has indicated the potential for stem cell therapies in treatment of conditions such as intervertebral disc degeneration, osteoarthritis and chronic tendinopathy (4, 5, 6, 8, 23). As such, stem cell therapies are at the forefront of research in regenerative medicine.

The regenerative potential of stem cell therapies has demonstrated immense promise in pain management techniques for musculoskeletal injuries and diseases, such as intervertebral disc degeneration. Specifically, ongoing clinical trials exploring the benefits of therapies containing umbilical cord-derived MSCs from Wharton's jelly region show encouraging therapeutic potential for reducing pain and inflammation and enhancing healing (2, 3).

Recent studies have demonstrated that MSCs derived from Wharton's Jelly may have the potential to reduce the apoptosis of intervertebral disc cells, called nucleus pulposus cells

(NPCs), that results from disc compression (4). Further, co-culturing MSCs derived from Wharton's Jelly with nucleus pulposus cells from degenerative intervertebral disc cells can induce MSC differentiation into NPCs and can activate favorable gene expression in degenerative NPCs (5). Clinically, MSCs derived from Wharton's jelly have been utilized in treatments aimed at repairing spinal cord, liver and heart tissue injuries and in the treatment of immune-mediated diseases (6).

Wharton's jelly is a gelatinous substance obtained from the connective tissue of human umbilical cords. In the umbilical cord, Wharton's jelly has several functions: to provide structural and protective support of the umbilical blood vessels and to provide a developing fetus with oxygen, glucose and amino acids while facilitating removal of waste products. Clinically, MSCs derived from Wharton's jelly may have the ability to stimulate cellular regeneration while eliciting little host immune response, as well as reducing pain and promoting healing through secretion of factors involved in immunomodulation. Regenerative medicine literature suggests the distinctive composition of Wharton's jelly may play an important role in its anti-inflammatory, pain-reducing, and enhanced healing capabilities (7).

Although MSCs can be isolated from myriad tissue types such as bone marrow and connective tissue of the umbilical cord, and are able to differentiate into osteoblasts, chondrocytes, adipocytes, and supported hematopoietic stroma, Wharton's jelly has been shown to contain the highest concentration of harvestable MSCs of allogenic tissues (2, 8). MSCs derived from umbilical cord Wharton's jelly region secrete factors thought to be involved in the regenerative and immunomodulatory abilities of MSCs (9). Further, the MSCs found in Wharton's jelly may also secrete factors that improve wound healing through enhanced intracellular communication (9).

In addition to the high concentration of MSCs, Wharton's jelly contains significant amounts of growth factors, cytokines, hyaluronic acid and exosomes (7). These constituents directly impact signaling pathways that contribute to the downstream anti-inflammatory, pain-reducing and enhanced healing effects of Wharton's jelly. Notably, recent research has demonstrated that decellularized Wharton's Jelly matrix (i.e. excluding MSCs) can promote cell differentiation of degenerative intervertebral disc cells towards a discogenic phenotype and can improve the phenotype of degenerative intervertebral disc cells (8). Further, decellularized Wharton's Jelly matrix was shown to maintain the viability of these discogenic cells (8). These remarkable findings demonstrate the regenerative capacity of Wharton's Jelly for treatment of intervertebral disc injuries (8).

Important growth factors found in Wharton's jelly include insulin-like growth factor-1 (IGF-1) carrier proteins that are known to improve osteogenic differentiation and induce MSCs into cell lines that develop into cartilaginous tissues, growth factors which are ligands for epidermal growth factor receptor (EGFR) and thus encourage growth and survival of osteogenic cells, and signaling proteins which are involved in bone remodeling called vascular endothelial growth factor (VEGF) (7, 11). The presence of such growth factors suggests the potential role of Wharton's jelly in regenerative medicine and is specifically applicable to designing pain

management therapies to repair damaged tissues without the need for surgery. The growth factors found in Wharton's jelly may be involved in modulating cellular mechanisms that combat chronic pain, joint and spine degeneration, and other musculoskeletal conditions due to their orthopedic-related nature. At present, many clinical and preclinical studies are investigating the potential of Wharton's jelly for treatment of various orthopedic conditions (12).

Additionally, many immunomodulatory cytokines found in Wharton's jelly are relevant to the field of regenerative medicine. Cytokines are substances that are produced by cells of the immune system and are often classified as either pro-inflammatory cytokines or anti-inflammatory cytokines. As the classification suggests, anti-inflammatory cytokines play critical roles in inhibiting inflammation, preventing hyperalgesia, and promoting wound healing. Important anti-inflammatory cytokines found in Wharton's jelly include chemokine ligand 5 (CCL5) which is associated with downstream signaling that leads to increased bone formation and a promoter of osteogenic differentiation called interleukin 6 receptor (IL-6R) (7, 13, 14). Pro-inflammatory cytokines are often associated with harmful inflammatory responses, but studies have also indicated that pro-inflammatory cytokines may play a role in initiating regenerative function of osteoclasts (7, 15). Such pro-inflammatory cytokines, called macrophage colony-stimulating factor (MCSF) and macrophage stimulating protein 1-alpha (MIP1- $\alpha$ ), are found in Wharton's jelly (7).

Furthermore, studies have demonstrated that Wharton's jelly contains significant amounts of hyaluronic acid (HA) (7). HA is found in many areas of the body and is relevant to various aging and musculoskeletal disorders, such as epidermal atrophy and knee osteoarthritis. Specifically, HA is an important component of synovial fluid, which acts as a lubricant in joints, and is produced by connective tissue cells such as chondrocytes and fibroblasts. As many musculoskeletal diseases like osteoarthritis are associated with reduced HA concentrations, conventional treatment of these diseases often includes HA injections called viscosupplementation injections (16). Recent literature has suggested that the combination of components in Wharton's jelly may provide improved efficacy as a treatment for knee osteoarthritis compared to HA injections alone, and currently there are clinical trials investigating this hypothesis (17).

In addition, the significant levels of extracellular vesicles including exosomes found in Wharton's jelly provides further insight into the potential therapeutic advantages of Wharton's jelly in regenerative medicine (7). Extracellular vesicles are essential to intercellular communication, but their functions extend far beyond this purpose. Extracellular vesicles relate to regenerative medicine through their role in MSC paracrine signaling and through their ability to release specific RNA species which can alter gene expression by modulating the epigenetic landscapes of target cells (18). Exosomes are a type of extracellular vesicle secreted from most stem cells and have been found to be significant secretory products of Wharton's jelly-derived MSCs. Studies suggest that the extracellular vesicles derived from Wharton's jelly show promising therapeutic avenues for numerous conditions ranging from myocardial ischemia/reperfusion injury to chronic kidney disease to knee osteoarthritis (10, 19, 20, 21)

Regenerative medicine research and preliminary clinical study data suggests that the amalgamation of the varied composition of Wharton's jelly, including the high concentration of MSCs and the presence of growth factors, cytokines, hyaluronic acid, and extracellular vesicles including exosomes, shows promising evidence that Wharton's jelly may exert anti-inflammatory, pain-reducing and enhanced healing effects in many diseases and conditions.

Notably, the MSCs found in umbilical cord-derived Wharton's jelly entail fewer ethical and safety concerns compared to stem cells isolated from adult tissue because they are harvested after birth, have a reduced likelihood of recipient rejection due to improved immunoprivilege, and demonstrate enhanced proliferation and maintenance of multipotency (5, 22).

Prior to the recent interest in regenerative medicine, the treatment of many musculoskeletal diseases relied heavily on surgical intervention. Although minimally invasive injections and procedures have become more common, these practices are not cures. As the field of regenerative medicine continues to grow, it is essential to consider that developing regenerative therapeutics for musculoskeletal diseases, such as MSCs derived from Wharton's jelly, may serve as a promising and non-surgical alternative to conventional treatments.

#### **Works Cited**

- (1) Roufosse, C A et al. "Circulating mesenchymal stem cells." *The international journal of biochemistry & cell biology* vol. 36,4 (2004): 585-97. doi:10.1016/j.biocel.2003.10.007
- (2) Friedenstein, A J et al. "Heterotopic of bone marrow. Analysis of precursor cells for osteogenic and hematopoietic tissues." *Transplantation* vol. 6,2 (1968): 230-47.
- (3) Marino, Luigi et al. "Mesenchymal Stem Cells from the Wharton's Jelly of the Human Umbilical Cord: Biological Properties and Therapeutic Potential." *International journal of stem cells* vol. 12,2 (2019): 218-226. doi:10.15283/ijsc18034
- (4) Zhao, Y-T et al. "Wharton's Jelly-derived mesenchymal stem cells suppress apoptosis of nucleus pulposus cells in intervertebral disc degeneration via Wnt pathway." *European review for medical and pharmacological sciences* vol. 24,19 (2020): 9807-9814. doi:10.26355/eurrev\_202010\_23190
- (5) Han, Zhihua et al. "Human Wharton's Jelly Cells Activate Degenerative Nucleus Pulposus Cells In Vitro." *Tissue engineering. Part A* vol. 24,13-14 (2018): 1035-1043. doi:10.1089/ten.TEA.2017.0340
- (6) Liao, L L et al. "Characteristics and clinical applications of Wharton's jelly-derived mesenchymal stromal cells." *Current research in translational medicine* vol. 68,1 (2020): 5-16. doi:10.1016/j.retram.2019.09.001

- (7) Gupta, Ashim et al. "Umbilical cord-derived Wharton's jelly for regenerative medicine applications." *Journal of orthopaedic surgery and research* vol. 15,1 49. 13 Feb. 2020, doi:10.1186/s13018-020-1553-7
- (8) Penolazzi, Letizia et al. "Extracellular Matrix From Decellularized Wharton's Jelly Improves the Behavior of Cells From Degenerated Intervertebral Disc." *Frontiers in bioengineering and biotechnology* vol. 8 262. 27 Mar. 2020, doi:10.3389/fbioe.2020.00262
- (9) Vangsness, C Thomas Jr et al. "Umbilical Cord Tissue Offers the Greatest Number of Harvestable Mesenchymal Stem Cells for Research and Clinical Application: A Literature Review of Different Harvest Sites." *Arthroscopy : the journal of arthroscopic & related surgery : official publication of the Arthroscopy Association of North America and the International Arthroscopy Association* vol. 31,9 (2015): 1836-43. doi:10.1016/j.arthro.2015.03.014
- (10) Bakhtyar, Nazihah et al. "Exosomes from acellular Wharton's jelly of the human umbilical cord promotes skin wound healing." *Stem cell research & therapy* vol. 9,1 193. 13 Jul. 2018, doi:10.1186/s13287-018-0921-2
- (11) Fortier, Lisa A et al. "The role of growth factors in cartilage repair." *Clinical orthopaedics and related research* vol. 469,10 (2011): 2706-15. doi:10.1007/s11999-011-1857-3
- (12) Main, Benjamin J et al. "Umbilical cord-derived Wharton's jelly for regenerative medicine applications in orthopedic surgery: a systematic review protocol." *Journal of orthopaedic surgery and research* vol. 15,1 527. 11 Nov. 2020, doi:10.1186/s13018-020-02067-w
- (13) Córdova, Luis A et al. "CCL2, CCL5, and IGF-1 participate in the immunomodulation of osteogenesis during M1/M2 transition in vitro." *Journal of biomedical materials research. Part A* vol. 105,11 (2017): 3069-3076. doi:10.1002/jbm.a.36166
- (14) Xie, Zhongyu et al. "Interleukin-6/interleukin-6 receptor complex promotes osteogenic differentiation of bone marrow-derived mesenchymal stem cells." *Stem cell research & therapy* vol. 9,1 13. 22 Jan. 2018, doi:10.1186/s13287-017-0766-0

- (15) Spiller, Kara L et al. "Sequential delivery of immunomodulatory cytokines to facilitate the M1-to-M2 transition of macrophages and enhance vascularization of bone scaffolds." *Biomaterials* vol. 37 (2015): 194-207. doi:10.1016/j.biomaterials.2014.10.017
- (16) Kosiński, Jakub et al. "HYALURONIC ACID IN ORTHOPEDICS." *Wiadomości lekarskie (Warsaw, Poland : 1960)* vol. 73,9 cz. 1 (2020): 1878-1881
- (17) Gupta, Ashim et al. "Safety and efficacy of umbilical cord-derived Wharton's jelly compared to hyaluronic acid and saline for knee osteoarthritis: study protocol for a randomized, controlled, single-blind, multi-center trial." *Journal of orthopaedic surgery and research* vol. 16,1 352. 31 May. 2021, doi:10.1186/s13018-021-02475-6
- (18) Matei, Andreea C et al. "Extracellular Vesicles as a Potential Therapy for Neonatal Conditions: State of the Art and Challenges in Clinical Translation." *Pharmaceutics* vol. 11,8 404. 11 Aug. 2019, doi:10.3390/pharmaceutics11080404
- (19) Lai, Ruenn Chai et al. "Exosome secreted by MSC reduces myocardial ischemia/reperfusion injury." *Stem cell research* vol. 4,3 (2010): 214-22. doi:10.1016/j.scr.2009.12.003
- (20) van Koppen, Arianne et al. "Human embryonic mesenchymal stem cell-derived conditioned medium rescues kidney function in rats with established chronic kidney disease." *PloS one* vol. 7,6 (2012): e38746. doi:10.1371/journal.pone.0038746
- (21) Song, Jun-Seob et al. "Implantation of allogenic umbilical cord blood-derived mesenchymal stem cells improves knee osteoarthritis outcomes: Two-year follow-up." *Regenerative therapy* vol. 14 32-39. 14 Jan. 2020, doi:10.1016/j.reth.2019.10.003
- (22) Millán-Rivero, Jose E et al. "Human Wharton's jelly mesenchymal stem cells protect axotomized rat retinal ganglion cells via secretion of anti-inflammatory and neurotrophic factors." *Scientific reports* vol. 8,1 16299. 2 Nov. 2018, doi:10.1038/s41598-018-34527-z
- (23) Lee, Sang Yoon et al. "Treatment of Lateral Epicondylitis by Using Allogeneic Adipose-Derived Mesenchymal Stem Cells: A Pilot Study." *Stem cells (Dayton, Ohio)* vol. 33,10 (2015): 2995-3005. doi:10.1002/stem.2110